PROCEEDINGS OF SCIENCE

# CP violation and Kaon weak matrix elements from Lattice QCD

arXiv:1512.02440v1 [hep-lat] 8 Dec 2015

**Nicolas Garron**[*†]
*Centre for Mathematical Sciences. School of Computing, Electronics and Mathematics, Plymouth University, Plymouth, PL4 8AA, United Kingdom*
*E-mail:* nicolas.garron@plymouth.ac.uk

In this short review, I present the recent lattice computations of kaon weak matrix elements relevant to $K \to \pi\pi$ decays and neutral kaon mixing. These matrix elements are key to the theoretical determination of the CP violation parameters $\varepsilon$ and $\varepsilon'$. Impressive progress have been achieved recently, in particular the first realistic computation of $\varepsilon'/\varepsilon$ with physical kinematics has been reported in [1]. The novelty is the $\Delta I = 1/2$ channel, whereas the $\Delta I = 3/2$ contribution is now computed at several values of the lattice spacing and extrapolated to the continuum limit. I will also present the status of $B_K$ and discuss its error budget, with a particular emphasis on the perturbative error. Finally I will review the matrix elements of neutral kaon mixing beyond the standard model and will argue that the discrepancy observed by different collaborations could be explained by the renormalisation procedure of the relevant four-quark operators.

*The 8th International Workshop on Chiral Dynamics*
*29 June 2015 - 03 July 2015*
*Pisa, Italy*

[*]Speaker.
[†]Supported by the Leverhulme Research grant RPG-2014-118

© Copyright owned by the author(s) under the terms of the Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International License (CC BY-NC-ND 4.0).

## 1. Introduction

Kaon physics has played and continues to play a central role in particle physics: CP violation, precision tests of the standard model, constraints on beyond-the-standard-model theories, etc. The recent progress achieved on the theoretical side, and in particular by the lattice community are improving drastically our theoretical understanding and open the door to new phenomenological studies. I review here the lattice computations related to $K \to \pi\pi$ decays and neutral kaon mixing. For more general recent reviews on lattice flavour physics see for example [2, 3].

## 2. $K \to \pi\pi$ decays and Lattice QCD

Various nice reviews are available on the subject, see for example [4]. Let me just recollect some basic facts about $K \to \pi\pi$ phenomenology, Assuming isospin symmetry, the decays $K \to \pi\pi$ can be written in terms of the amplitudes

$$A\left[K \to (\pi\pi)_I\right] = A_I e^{i\delta_I} \tag{2.1}$$

where $I$ denotes the isospin of the two-pion state, either 0 or 2, and $\delta_I$ is the corresponding strong phases. The parameters of indirect (resp. direct) CP violation, $\varepsilon$ (resp. $\varepsilon'$) are given by

$$\varepsilon = \frac{A\left[K_L \to (\pi\pi)_0\right]}{A\left[K_S \to (\pi\pi)_0\right]} . \tag{2.2}$$

$$\frac{\varepsilon'}{\varepsilon} = \frac{1}{\sqrt{2}}\left(\frac{A\left[K_L \to (\pi\pi)_2\right]}{A\left[K_L \to (\pi\pi)_0\right]} - \frac{A\left[K_S \to (\pi\pi)_2\right]}{A\left[K_S \to (\pi\pi)_0\right]}\right) \tag{2.3}$$

The first measurement of $\varepsilon$ is the well-known discovery of indirect CP violation due to Christenson, Cronin, Fitch and Turlay [5] in 1964, for which Cronin and Fitch were awarded a Nobel prize. $\varepsilon'$ has a long experimental history as it took a tremendous effort to measure direct CP violation. The final measurements are due to NA48 at Fermilab and KTeV at CERN [6, 7], the averages read

$$|\varepsilon| = 2.228(11) \times 10^{-3} , \tag{2.4}$$

$$Re\left(\frac{\varepsilon'}{\varepsilon}\right) = 16.6(2.3) \times 10^{-4} . \tag{2.5}$$

In a theoretical approach, the standard framework to study $K \to \pi\pi$ decay is the $\Delta S = 1$ effective Hamiltonian obtained after integrating out the heavy degrees of freedom. In the three-flavour theory, it reads (see for example [8], [9])

$$H_W = \frac{G_F}{\sqrt{2}} V_{us}^* V_{ud} \sum_{i=1}^{10} \left[z_i(\mu) + \tau y_i(\mu)\right] Q_i(\mu) , \tag{2.6}$$

where $G_F$ is the Fermi constant. The short-distance effects, which can be computed in perturbation theory are factorised into the so-called Wilson coefficients, $y_i, z_i$ whose expression can be found in [8]. $V_{ij}$ are CKM matrix elements, $\tau = V_{ts}^* V_{td} / V_{us}^* V_{ud}$ and $\mu$ is an energy scale which can be thought as a cut-off. The four-quark operators $Q_i$ can be obtained from Feynman diagrams and after operator-product-expansion they can be expressed as

Current-Current:

$$Q_1 = (\bar{s}^i\gamma_\mu(1-\gamma_5)d^i)(\bar{u}^j\gamma_\mu(1-\gamma_5)u^j) \tag{2.7}$$
$$Q_2 = (\bar{s}^i\gamma_\mu(1-\gamma_5)d^j)(\bar{u}^j\gamma_\mu(1-\gamma_5)u^i) \tag{2.8}$$

QCD Penguins:

$$Q_3 = (\bar{s}^i\gamma_\mu(1-\gamma_5)d^i)\sum_{q=u,d,s}(\bar{q}^j\gamma_\mu(1-\gamma_5)q^j) \tag{2.9}$$
$$Q_4 = (\bar{s}^i\gamma_\mu(1-\gamma_5)d^j)\sum_{q=u,d,s}(\bar{q}^j\gamma_\mu(1-\gamma_5)q^i) \tag{2.10}$$
$$Q_5 = (\bar{s}^i\gamma_\mu(1-\gamma_5)d^i)\sum_{q=u,d,s}(\bar{q}^j\gamma_\mu(1+\gamma_5)q^j) \tag{2.11}$$
$$Q_6 = (\bar{s}^i\gamma_\mu(1-\gamma_5)d^j)\sum_{q=u,d,s}(\bar{q}^j\gamma_\mu(1+\gamma_5)q^i) \tag{2.12}$$

EW Penguins:

$$Q_7 = (\bar{s}^i\gamma_\mu(1-\gamma_5)d^i)\sum_{q=u,d,s}e_q(\bar{q}^j\gamma_\mu(1+\gamma_5)q^j) \tag{2.13}$$
$$Q_8 = (\bar{s}^i\gamma_\mu(1-\gamma_5)d^j)\sum_{q=u,d,s}e_q(\bar{q}^j\gamma_\mu(1+\gamma_5)q^i) \tag{2.14}$$
$$Q_9 = (\bar{s}^i\gamma_\mu(1-\gamma_5)d^i)\sum_{q=u,d,s}e_q(\bar{q}^j\gamma_\mu(1-\gamma_5)q^j) \tag{2.15}$$
$$Q_{10} = (\bar{s}^i\gamma_\mu(1-\gamma_5)d^j)\sum_{q=u,d,s}e_q(\bar{q}^j\gamma_\mu(1-\gamma_5)q^i) \tag{2.16}$$

The matrix element of these four-quark operators capture the strong dynamics of the theory. We have neglected the operators which emerge from the electric and magnetic dipole part of the electromagnetic and QCD penguins. (See the talk by V.Lubicz at Lattice'14 and [10] for a recent lattice study by the ETM collaboration.) These 10 operators do not form a basis of the $\Delta S = 1$ four-quark operators in four dimensions, as they are not linearly independent. Following [11], we build a 7-operators basis:

$$Q'_1 = 3Q_1 + 2Q_2 - Q_3 \tag{2.17}$$
$$Q'_2 = \frac{1}{5}(2Q_1 - 2Q_2 + Q_3) \tag{2.18}$$
$$Q'_3 = \frac{1}{5}(-3Q_1 + 3Q_2 + Q_3) \tag{2.19}$$
$$Q'_i = Q_i\ , i \in \{5,6,7,8\}\ . \tag{2.20}$$

The $Q'_i$ falls into three different irreducible representations of $SU_L(3)\times SU_R(3)$: $Q'_1$ transforms as a $(27,1)$, the QCD penguins[1] $Q'_{2,3,5,6}$ as $(8,1)$ and the QED pengiuns as $(8,8)$.

Obtaining a reliable evaluation of the matrix elements $\langle\pi\pi|Q'_i|K\rangle$ is the most difficult part of the computation. Since one needs a non-pertubative framework, lattice QCD is a natural candidate. In the last thirty years, many attempts have been made to evaluate these matrix elements, using either effective theories or lattice simulations (or combinations of both), se for example [12, 13, 14, 15, 16, 17, 18] and reference therein.

[1] $Q'_{2,3}$ are actually combinations of current-current and QCD penguin operators.

From the lattice point of view, the first difficulty is to simulate the kinematic situation, in particular the final state made of two hadrons with non-vanishing momenta. This problem was formalised in 1990 by Maiani and Testa who showed that the physical amplitudes could not be extracted from "standard" eulcidean lattice simulations [19]. An alternative based on $\chi$PT was proposed in [20]: the matrix elements of interests can be obtained from those of $K \to \pi$ and $K \to$ vacuum, which are numerically much simpler. This indirect approach was first used for while, see for example [21, 22, 11]. However the conclusion of the extensive quenched studies [22, 11] is rather negative: extracting the matrix elements with a fully controlled error turned out to be very hard. One problem comes from the fact that $SU(3)$ $\chi$PT converges poorly at the kaon scale (see also [23]) [2].

What is now known as the Maiani-Testa no-go theorem was solved in an very elegant way by Lellouch and Lüscher in [26]. The crucial point is that in finite volume the spectrum is discrete, and the size of the box can be fine-tuned such that the pions will take the desired momentum.

## 3. The $\Delta I = 3/2$ channel

We first consider the amplitude of $K \to (\pi\pi)_{I=2}$ decays, there are several simplifications in this channel, most notably 1. there is no disconnected diagram, and 2. only three operators contribute. The first realistic computation (with dynamical quarks, physical kinematics and nearly-physical pion mass) was performed by the RBC-UKQCD collaborations [27, 28] with Domain-Wall fermions, a discretisation of the QCD lagrangian which preseves chiral-flavour symmetry almost exactly.

Although the method used in [27, 28] is based on the Lellouch-Lüscher approach, an important ingredient is the Wigner-Eckart theorem, which tells us that the matrix elements of interest are related to those of the unphysical process $K^+ \to \pi^+\pi^+$ (in the isospin limit). Using a peculiar choice boundary conditions, these matrix elements (with physical momenta) can be extracted using standard lattice methods. The first simulation was done at a single value of the lattice spacing ($a^{-1} \sim 1.375$ GeV, ie $a \sim 0.1435$ fm) on the so-called IDSDR lattice (ID) [29] with a pion mass of 140 MeV. (Strictly speaking this "physical pion" is partially quenched, the unitary pion mass was somewhat heavier: 170 MeV). In this work, the matrix elements are renormalised non-perturbatively , with the Rome-Southampton method [30]. Since this lattice spacing is rather coarse, the renormalisation is first performed at a rather low value of the momentum scale ($\mu \sim 1.1$ GeV). In a second step, the same renormalisation factors are evaluated on finer lattices (called Iwasaki (IW) lattices) and the (universal) continuum scale-evolution matrix to 3 GeV is obtained from [31], schematically:

$$Z^{ID}(3\,\mathrm{GeV}, a_{ID}) = \lim_{a_{IW}\to 0} \left[ Z^{IW}(3\,\mathrm{GeV}, a_{IW}) \left( Z^{IW}(1.1\,\mathrm{GeV}, a_{IW}) \right)^{-1} \right] Z^{ID}(1.1\,\mathrm{GeV}, a_{ID}) \,. \quad (3.1)$$

More recently, the RBC-UKQCD collaborations have reported on $2+1$ lattice QCD simulations with physical pion masses [29], which have been possible thanks to a new formulation of the Domain-Wall disctretisation [32]. These lattices have been used to improve on the determination of

[2] An interesting proposal based on Chiral-Scale Perturbation Theory has been presented at the conference by Lewis Tunstall, [24] see also [25].




**Figure 1:** Real and Imaginary part of $A_2 = A\left[K \to (\pi\pi)\right]_{I=2}$. The triangle represents the 2012 computation on the IDSDR and the blue points the 2014 determinations on the new ensembles (statistical error only), from which a continuum limit is extracted and shown in magenta (statistical and systematic errors combined). For the IDSDR points, we show both the statistical and the systematic error, largely dominated by the discretisation artefacts.

$A_2$: the main source of error was the discretisation effects, the new computation [33] involves two lattice spacings of $a \sim 0.011$ and $a \sim 0.084$ fm, reducing the systematic error by roughly a factor 2 for the real part and a factor 1.5 for the imaginary part. Thanks to these new lattice determinations, the current errors on the theoretical determination of $A_2$ are of the order of 10%.

## 4. Including the $\Delta I = 1/2$ channel

A complete determination of $A\left[K \to \pi\pi\right]_{I=0}$ has been a long-standing challenge for the lattice community. A first "pilot" computation with dynamical fermions was reported by RBC-UKQCD in 2011 [34]. This computation was unphysical in the sense that the amplitudes were computed at threshold and the quark masses were heavier that the physical ones, however all the required diagrams were determined (including the disconnected ones) showing the numerical feasibility of the approach. The main remaining difficulty was to implement the physical kinematics, ie the ability of extracting the matrix element of interests, with the pion states having the right momenta. The Wigner-Eckart/boundary condition trick cannot used in the $\Delta = 3/2$ channel cannot be used for the full computation as it violates isospin. Instead, the RBC-UKQCD collaboration have generated new ensemble with $G-$ parity boundary conditions [35, 36], as reported by Christopher Kelly in a plenary session of Lattice 2015 [37], see also the plenary review given by Andreas Jüttner at the same conference [38]. From a more technical point of view, this computation requires the evaluation of all-to-all propagators and noise reduction techniques. The results read

$$Re(A_0) = 4.66(1.00)(1.21) \times 10^{-7}\text{GeV} \tag{4.1}$$
$$Im(A_0) = -1.90(1.23)(1.04) \times 10^{-11}\text{GeV} \tag{4.2}$$

and the corresponding theoretical value for $\varepsilon'\varepsilon$

$$Re(\varepsilon'/\varepsilon) = 1.38(5.15)(4.43) \times 10^{-4}\,, \tag{4.3}$$

which is an approximate agreement($\sim 2.1\sigma$) with the experimental value $16.6(2.3)\times 10^{-4}$. Rather than concluding that a significant deviation of the Standard Model prediction has been found, we note that the error is much larger than experimental one. From a phenomenological point of view, at this level of precision, these results do not invalidate the standard model, neither do they rule the need for new-physics in $K\to\pi\pi$ decays. The important point is that for the first time $\varepsilon'/\varepsilon$ has been computed with a full error budget, all the different contributions of the seven linearly independent operators are computed with controlled errors and a precision which can be systematically improved. Now that the technology has been developed, reaching a precision of, say, 10% should be possible in the close future. In addition to reducing the statistical error, the simulation can be done on finer lattices and extrapolated to the continuum limit. Another systematic error is due to the truncation of the perturbation series (needed to compute the Wilson coefficients). The renormalisation was performed at a scale of $\mu\sim 1.5$ GeV in order to keep the discretisation effects under control. Clearly this can be improved by running non-perturbatively to a higher scale, as done for the $\Delta I=3/2$ channel. Reducing the theoretical error on the matrix elements of $O_i$ (and therefore on n $\varepsilon'/\varepsilon$ ) will provide a crucial test of the standard model, indeed we might actually see signs of new physics. It is also worth noting that another computation (done at threshold) has been done with Wilson fermions [39].

## 5. The $\Delta I=1/2$ rule

The "$\Delta I=1/2$ rule" refers to the fact that the $I=0$ channel is favoured over the $I=2$ channel by important factor $1/\omega$ defined by

$$\omega=\frac{A\left[K_S\to(\pi\pi)_2\right]}{A\left[K_S\to(\pi\pi)_0\right]} \tag{5.1}$$

Experimentally this number is of order $\omega\sim 1/22$ whereas one would naively expects $1/2$ [40, 41]. The question whether or not the remaining factor of $\sim 10$ can be explained entirely by some surprisingly large QCD effects has been a very-long standing puzzle. It also shows the need for a better understanding of the non-perturbative regime. Several attempts to study the $\Delta I=1/2$ rule on the lattice have been made. For example, an ongoing project based on the role of the charm quark has been developed in [42, 43, 44, 45], see also [46].

In 2013, the RBC-UKQCD collaborations reported on a study of the origin of this enhancement [47]. The amplitude $A_2$ was computed with physical kinematics whereas $A_0$ was computed at threshold. The real part of $A_2$ is largely dominated by single four-quark operator (the contributions of the electroweak penguins are negligible with respect to the tree-level diagram). This operator has a $(V-A)\times(V-A)$ Dirac structure and transform as a $(27,1)$. Two contractions ① and ② contribute, they differ by their colour structure, as shown in figure 2. The conventions are such that the real part of $A_2$ can be approximated by sum ① + ② which are shown in figure 2. The naive expectation is that ② $\sim\frac{1}{3}$①. However the observation made in [47] is that ② $\sim -0.7$①. Therefore, there is important cancellation in the numerator of Eq.(5.2) which is completely unexpected from the naive factorisation framework. Similarly the main contribution to $Re(A_0)$ is proportional




**Figure 2:** The dominating contribution to the real part of the amplitude $A_2$ of $K \to \pi\pi$ is proportional to sum of the two contractions ① and ②. The two contractions differ by their colour structure, as indicated by the colour indices $i$ and $j$. The label $L$ stands for the left-handed structure $\gamma_\mu(1-\gamma_5)$.

to $2① - ②$. Hence, the aforementioned relative sign between ① and ② also contributes to enhancement in the denominator of $\omega$ (compared to the naive expectation).

$$\omega \sim \frac{Re(A_2)}{Re(A_0)} \sim \frac{① + ②}{2① - ②} \tag{5.2}$$

The two recent lattice computations, the threshold one [39] and the one with physical kinematics [1] also observe this sign difference, which seems to be at the origin of the $\Delta I = 1/2$ rule. However, in order to confirm that the $\Delta I = 1/2$ effect is a pure non-perturbative QCD effect, a little bit of patience is required as the precision on $A_0$ has to be improved. The theoretical error affecting the amplitudes is expected to decrease by a factor two in the next couple of years, it is very likely that we will then have the answer to this question. Note that this sign also discussed in [4, 48], see also. [17, 8, 9].

## 6. Neutral kaon mixing and indirect CP violation in the Standard Model

In the Standard Model picture, neutral kaon mixing is dominated by $W$-exchange box diagrams as illustrated in figure 3, a well-known loop-suppressed flavour changing neutral current. By performing an operator product expansion, one can factorise the long-distance effects into the




**Figure 3:** Box diagram contributing to $K^0 - \bar{K}^0$ mixing in the SM.

matrix element of a four quark operator:

$$\langle \bar{K}^0 | O_1^{\Delta S=2} | K^0 \rangle = \langle \bar{K}^0 | (\bar{s}_i \gamma_\mu (1-\gamma_5) d_i) (\bar{s}_j \gamma_\mu (1-\gamma_5) d_j) | K^0 \rangle \ . \tag{6.1}$$

Clearly, because of the W-exchange, the Dirac structure is "(Vector-Axial) $\times$ (Vector-Axial) ". It is quite remarkable that only one four-quark operator contribute: even turning on QCD does not add

any Dirac structure because the operator given in eq. 6.1 is invariant under Fierz re-arrangement. In a (continuum) massless renormalisation scheme, it does not mix with other four-quark operators, nor with lower dimensional operators. It is also the case on the lattice if chiral symmetry is preserved.

Once the considered matrix element has been computed non-perturbatively using lattice techniques, its result is combined with the value of the Wilson coefficient $C(\mu)$ of continuum perturbation theory and experimental observables, such as the mass difference $\Delta_{M_K} = m_{K_L} - m_{K_S}$ and $\varepsilon_K$ to obtain important constraints on the CKM matrix elements. Schematically, one obtains

$$\varepsilon_K = C(\mu) \times \langle \bar{K}^0 | O_1^{\Delta S=2} | K^0 \rangle(\mu) \times \mathscr{F}(V_{ij}^{CKM}, m_K, f_K, \Delta M_K, \ldots) \,, \tag{6.2}$$

where $\mathscr{F}$ is a known function of the CKM factors and of well-measured quantities.

A convenient parametrisation of this operator is the well-known bag parameter $B_K$,

$$B_K(\mu) \equiv \frac{\langle \bar{K}^0 | O_1^{\Delta S=2} | K^0 \rangle(\mu)}{\frac{8}{3} f_K^2 m_K^2}. \tag{6.3}$$

where $f_{K^-} = 156.1$ MeV and $\mu$ is a renormalisation scale, usually 2 or 3 GeV. On the lattice, the bare bag parameter can be obtained from a ratio of correlators

$$r_{B_K}^{bare} = \frac{\langle P^\dagger(t_2) O_1^{\Delta S=2}(t_o) P^\dagger(t_1) \rangle}{\langle P^\dagger(t_2) A_0(t_o) \rangle \langle A_0(t_o) P^\dagger(t_1) \rangle} \tag{6.4}$$

where $P^\dagger$ create a light-strange pseudo-scalar particle (which would be a kaon in the continuum at the physical value of the quark masses) and $A_0$ is the time component of the corresponding axial current. Up to some numerical factors, the bare bag parameter $B_K^{bare}$ is obtained from fitting $r_{B_K}^{bare}$ in the asymptotic euclidian time region $t_2 \ll t_o \ll t_1$

$B_K$ is a standard lattice quantity, nowadays it is computed with an accuracy of a few percents [49][50][51]. The FLAG 2013 average for $N_f = 2+1$ is

$$\hat{B}_K \;=\; 0.7661(99) \,, \qquad N_f = 2+1 \,, \tag{6.5}$$

it is largely dominated by the BMWc result $\hat{B}_K = 0.773(8)_{stat}(3)_{syst}(8)_{PT}$. The other references are [52, 53, 54, 55, 50, 56, 57, 58, 59].

Let us consider the dominant sources of error: FLAG 2013 explains that the total error of $99 \sim 1.3\%$ can be roughly seen as the combination of 0.4% statistical and 1.2% systematic, mainly due to perturbation theory: $(33)_{stat} + (93)_{syst}$. Although $B_K$ is extracted and (in most cases) renormalised non-perturbatively on the lattice, perturbation theory is used to convert the result to the renormalisation-group-invariant (RGI) quantity $\hat{B}_K$, or alternatively to $\overline{\text{MS}}$. Naturally, different collaborations estimate the perturbative error in different ways, and this estimation is of course affected by some subjective judgement. Indeed this error changes by a factor two or three depending on the estimation. In its 2013 review, it seems that FLAG chose an uncertainty very close to the one quoted by BMW (1%), whereas RBC-UKQCD quoted an error of $\sim 2\%$, based on a multiple-scheme evaluation. Actually by changing the intermediate schemes, RBC-UKQCD find that the results change by 8% if the matching is done at $\mu = 3$GeV and by 12% if $\mu = 2$GeV. The current situation is illustrated on table 1, where we have collected the most recent computation of $B_K$.

| Collaboration | $N_f$ | Discretisation | Result |
|---|---|---|---|
| RBC-UKQCD [29] | $2+1$ | Domain-Wall | $0.5293(17)_{stat}(106)_{PT}$ |
| SWME [62] | $2+1$ | Staggered | $0.518(3)_{stat}(26)_{syst}$ |
| ETM[63] | $2+1+1$ | Twisted Mass | $0.506(17)_{stat+syst}(3)_{PT}$ |

**Table 1:** Collection of recent results for $B_K^{\overline{\text{MS}}}(3\text{GeV})$. In [29], the first error is statistical but is much larger than the systematic ones, except for the perturbative error which is written as $(PT)$. For the latter, it is interesting to note that RBC-UKQCD quotes an error of 2% whereas ETM quotes 0.6%. In [62] this error is not distinguished from other uncertainties due to the renormalisation procedure. BMWc result is not quoted here as it uses different convention, but quotes a perturbative error of 1%. This perturbative error is discussed in detail in the text.

The importance of this perturbative error can be made clear by looking at, for example, the result obtained by RBC-UKQCD [29]

$$\hat{B}_K = 0.7499(24)_{stat}(150)_{PT} \tag{6.6}$$

$$B_K^{(\not{q},\not{q})}(3\text{GeV}) = 0.5341(18)_{stat} \tag{6.7}$$

where the first error is statistical (however it is much larger than the other errors on $B_K^{bare}$) and the second error is the systematic error on the renormalisation, largely dominated by the perturbative matching. This contrasts with $B_K^{(\not{q},\not{q})}$ which is fully non pertutbative, renormalised in the SMOM $(\not{q},\not{q})$-scheme at the scale $\mu = 3$ GeV. Without this perturbative error, the error would be of around 0.3%.

In the future, the lattice community will probably have to find an agreement on how to estimate this uncertainty as it is the dominant one. Another way to improve the situation is obviously to reduce this perturbative error. One way would be to compute explicitly the next order in perturbation theory. The matching coefficient is currently known at next-to-leading order. Going further requires to determine the matching coefficient at the two-loop level (three-loop anomalous dimension). Alternatively, on could perform the matching at a higher scale; this could be achieved by computing the running non-perturbatively, for example using the Schrödinger functional [60] or a (S)MOM-scheme, as presented in [61].

## 7. Neutral kaon mixing Beyond the Standard Model

In the standard model operator, only the operator $(V-A)\times(V-A)$ contributes to kaon oscillations, because the process can only occur through $W$ exchanges. Beyond the standard model, we have to include new Dirac-colour structures, as for example both left-handed and right-handed currents can contribute to $K^0$-$\bar{K}^0$ mixing (and therefore to $\varepsilon_K$). Hence, in addition to $O_1$ introduced in eq.6.1, one introduces new $\Delta S = 2$ four-quark operators. A possibility (the so-called SUSY-basis) is [64, 65] [3]

$$O_2^{\Delta S=2} = (\bar{s}_i(1-\gamma_5)d_i)\,(\bar{s}_j(1-\gamma_5)d_j), \tag{7.1}$$

[3] An alternative basis is the one given by [66] .

$$O_3^{\Delta S=2} = (\bar{s}_i(1-\gamma_5)d_j)(\bar{s}_j(1-\gamma_5)d_i), \tag{7.2}$$
$$O_4^{\Delta S=2} = (\bar{s}_i(1-\gamma_5)d_i)(\bar{s}_j(1+\gamma_5)d_j), \tag{7.3}$$
$$O_5^{\Delta S=2} = (\bar{s}_i(1-\gamma_5)d_j)(\bar{s}_j(1+\gamma_5)d_i). \tag{7.4}$$

The BSM operators $O_{i\geq 2}^{\Delta S=2}$ mix pair-wise: $O_2^{\Delta S=2}$ with $O_3^{\Delta S=2}$ (in the chiral limit they transform under a $(6,\bar{6})$ irreducible representation of $SU(3)_L \times SU(3)_R$) and $O_4^{\Delta S=2}$ with $O_5^{\Delta S=2}$ (the corresponding group irrep being $(8,8)$). These four-quark operators appear in the generic effective $\Delta S = 2$ Hamiltonian

$$H^{\Delta S=2} = \sum_{i=1}^{5} C_i(\mu)\, O_i^{\Delta S=2}(\mu) + , \sum_{i=1}^{3} \tilde{C}_i(\mu) \tag{7.5}$$

where the Wilson coefficient $C_i(\mu), \tilde{C}_i(\mu)$ depend on the details of the new-physics model under consideration but the matrix elements $\langle \bar{K}^0 | O_i^{\Delta S=2} | K^0 \rangle$ are model independent. The operators $\tilde{O}_{1,2,3}$ are obtained from $O_{1,2,3}$ by replacing $(1-\gamma_5)$ by $(1+\gamma_5)$. In QCD with parity conserved, these operators are redundant and therefore discarded in the following.

*A priori*, one would expect that the relevant matrix elements $\langle \bar{K}^0 | O_1^{\Delta S=2} | K^0 \rangle$ can be obtained with an accuracy comparable to the one of the Standard Model one. However only few studies of the full set of BSM operators have been published and the history is quite interesting. First of all, in the quenched approximation, the results from [67] obtained with Ginsparg-Wilson fermion (which exhibit an exact chiral-flavour symmetry) and non-perturbative renormalisation were very different from the previous study, done with tree-level $O(a)$-improved Wilson fermions [68]. The difference was attributed to the renormalisation.

The first computation performed with dynamical fermions was reported by RBC-UKQCD [69] in 2012 and was done with $N_f = 2+1$ Domain-Wall fermions. It was followed shortly by a $N_f = 2$ twisted-mass computation of the ETM collaboration, done with several lattice spacings [70] and these two first computations are in reasonable agreement (slightly more than2% in the worse case) In 2013, the SWME collaboration reported their results, obtained with $N_f = 2+1$ flavours of improved staggered fermions [71]. A noticeable disagreement with the previous studies was found for two of the matrix elements ($O_4$ and $O_5$ of the SUSY basis). Very recently, the ETM collaboration have repeated their computation with $N_f = 2+1+1$ flavours (using again twisted mass QCD), and essentially confirmed their $N_f = 2$ results [63] (although for $B_5$ the agreement is only within $\sim 3\sigma$). Even more recently, SWME have extended their study by adding more ensembles, [62] improving the extrapolation to the physical point, and they confirmed the disagreement with the other collaboration. Since the results have been extrapolated to the continuum limit, one does not expect that the discretisation used (Domain-Wall, Twisted-Mass, or Staggered) is responsible of the discrepancy. A first suspect is the renormalisation procedure. ETM and RBC-UKQCD employ a non-perturbative method, based on the Rome-Southampton method[30], whereas SWME uses a perturbative matching. Another possibility could an underestimated systematic error due to a chiral log. Although this sounds more unlikely it could be investigated, in a first step by extrapolating quantities which have different chiral behaviour (the $B's$, the $R's$ and the $G's$) and comparing the results. Eventually these quantities will be computed directly at the physical value of the pion mass, as done for the Standard Model contribution.

**Normalisation.** The matrix elements of these four-quark operators are usually given in terms of the so-called Bag-parameters,

$$B_i(\mu) = \frac{\langle \bar{K}^0 | O_i(\mu) | K^0 \rangle}{\langle \bar{K}^0 | O_i(\mu) | K^0 \rangle_{VS}}, \tag{7.6}$$

where $VS$ is the vacuum-saturation approximation. On the lattice, this is by achieved by computing a ratio of three-point function over a product of two-point functions, such as $r_{B_K}^{bare}$ defined in eq. (6.4). In the case of the Standard Model operator, the denominator is known in terms of physical quantities $f_K$ and $m_K$, as shown in eq. (6.3). This normalisation is convenient because the bags parameters are dimensionless, the numerator and the denominator are very similar, therefore systematic errors are likely to cancel out in the ratio, and because the denominator is known in terms of physical quantities. However for the BSM operators, the corresponding vaccum saturation involve matrix element of the pseudo-scalar density, and on the lattice the corresponding $r_{B_{i>1}}^{bare}$ usually contains the product of pseudo-scalar pseudo-scalar two-point function $\langle P^\dagger(t_2)P(t_o)\rangle\langle P(t_o)P^\dagger(t_1)\rangle$, which is approximated by $m_K^4 f_K^2/(m_s+m_d)^2$. First, this is only an approximation, but also the fact that quark masses appear imply new ambiguities such as scale and scheme dependencies. Numerically, we also find the ratio of three-point functions is better determined than the ratio three-point over two point functions.

This problem is well known and several alternatives were proposed in the literature, see for example [68]. Let us mention explicitly the solution proposed in [67]. Denoting by $P$ the simulated pseudo-scalar particle (kaon) of mass $m_P$ and decay constant $f_P$, the $R$'s are defined by

$$\mathscr{R}_i\left(\frac{m_P^2}{f_P^2}, \mu, a^2\right) = \left[\frac{f_K^2}{m_K^2}\right]_{\text{expt}} \left[\frac{m_P^2}{f_P^2}\frac{\langle \bar{P}|O_i(\mu)|P\rangle}{\langle \bar{P}|O_1(\mu)|P\rangle}\right]_{\text{latt}}. \tag{7.7}$$

such that at the physical point $m_P = m_K = m_K^{\text{expt}}$ and $a = 0$

$$R_i(\mu) = \mathscr{R}_i\left(\frac{m_K^2}{f_K^2}, \mu, 0\right) = \frac{\langle \bar{K}^0|O_i(\mu)|K^0\rangle}{\langle \bar{K}^0|O_1(\mu)|K^0\rangle} \tag{7.8}$$

is directly proportional to the ratio of a BSM contribution to the SM one.

Another possibility, advocated for example in [71], is to define products and ratios of bag parameters (called $G$) such that the leading chiral logarithms cancel out. This cancellation occurs at best at every order of the chiral expansion, or in the worst case to next-to-next-to-leading order (NNLO). Such quantities were introduced in [72] for $SU(3)$ chiralperturbation theory and expanded later in the context of $SU(2)$ staggered chiral perturbation theory in [73]. The problem of the normalisation ambiguity is absent for the ratios, but still there for the product. However the advantage is that the chiral extrapolations are hugely simplified. Having different normalisations and fit Ansätze help to a better control of the systematic error. One can for example extract the $B$ directly with some chiral logs, of reconstruct them from the linearly extrapolated $G$.

**New RBC-UKQCD results (preliminary) and tentative explanation of the disagreement.**
I am presenting now a work in progress with the RBC-UKQCD collaborations, preliminary results have been presented by R.J.Hudspith at Lattice 2015 [74].

| $R^{\text{SUSY}}(3\,\text{GeV})$ | SMOM $-\gamma_\mu$ | $\overline{\text{MS}}$ |
|---|---|---|
| $R_2$ | $-19.11(43)(19)(25)$ | $-19.48(44)(20)(25)$ |
| $R_3$ | $5.76(14)(15)(07)$ | $6.08(15)(16)(08)$ |
| $R_4$ | $40.1(08)(17)(09)$ | $43.1(09)(18)(10)$ |
| $R_5$ | $11.13(21)(79)(25)$ | $10.99(20)(78)(25)$ |

**Table 2:** RBC-UKQCD preliminary results for the BSM $\Delta S = 2$ ratio $R$ in the SUSY basis, in the $\gamma_\mu$-SMOM scheme and in $\overline{\text{MS}}$ at 3 GeV. The quantities $R$ are the ratios of the BSM matrix elements over the SM contributions. Errors are statistics, discretisation, chiral, respectively.

**Possible explanation.** In order to understand the source of the disagreement, we have extended our previous study [69] in several ways, most notably 1. by adding another lattice spacing 2. by investigating new non-perturbative renormalisation (NPR) SMOM schemes, in the spirit of the schemes introduced in [75]. Our main results are presented in Table 2, in a SMOM scheme, and in $\overline{\text{MS}}$ (we would like to thank Christoph Lehner for computing the conversion factors for $B_2$ and $B_3$). A comparison of results for the bag parameters can be found in Table 7. Although our error budget is not complete yet, we find that if we use the standard RI-MOM scheme proposed in [30] and match to the $\overline{\text{MS}}$ scheme defined in [66], our results are in a decent agreement with ETMc. Surprisingly enough. if we use a SMOM scheme, our results are much closer to the results quoted by SWME, for which the renormalisation is performed perturbatively. The SMOM schemes are known to be superior to standard RI-MOM schemes: they behave better non-perturbatively in the infrared (the pion pole contamination is suppressed because of the absence of exceptional channel) and perturbatively. Our suspicion is that the procedure employed to remove the pion pole contamination (needed in the RI-MOM case but absent for the SMOM schemes) could also affect the ultraviolet behaviour, see for example [76]. The systematic errors associated with this procedure are very hard to estimate, they could have been underestimated. We do not advocate to use the bag parameters for our central values, we only show them in order to compare with collaborations. We find that the $R$'s have smaller systematic and give much more reliable results. On the negative side, we find that - regardless of the normalisation - the discretisation effects are larger than expected. Although these artefact are moderate (we quote 7% from the $a^2$ slope in the worse case) they are larger than what is usually found with Domain-Wall fermions. In the future, we are planning to add a third lattice spacing in order in order to have a better handle on these lattice artefacts.

## 8. Conclusions and outlook

Kaon physics is experiencing a new youth (see for example [77]); to a great extent this is due to the impressive progress achieved recently by the lattice community. The computation of $K \to (\pi\pi)_{I=2}$ is reaching a mature stage and a first computation $K \to (\pi\pi)_{I=0}$ with physical kinematic and complete error budget has recently been reported by the RBC-UKQCD collaboration. The results of these computations have a important role to play in particle physics phenomenol-

| | ETM′12 | ETM′15 | RBC – UKQCD′12 | SWME′15 | RBC – UKQCD′15(prelim.) | |
|---|---|---|---|---|---|---|
| *interm.* | | | | | | |
| *scheme* | $RI-MOM$ | $RI-MOM$ | $RI-MOM$ | $1-loop$ | $RI-SMOM$ | $RI-MOM$ |
| $B_2$ | 0.47(2) | 0.46(3) | 0.43(5) | 0.525(1)(23) | 0.488(7)(17)(2) | 0.417(6)(2)(2) |
| $B_3$ | 0.78(4) | 0.79(5) | 0.75(9) | 0.772(5)(35) | 0.743(14)(64)(3) | 0.655(12)(44)(2) |
| $B_4$ | 0.75(3) | 0.78(5) | 0.69(7) | 0.981(3)(61) | 0.920(12)(12)(4) | 0.745(9)(28)(3) |
| $B_5$ | 0.60(3) | 0.49(4) | 0.47(6) | 0.751(8)(68) | 0.707(8)(34)(3) | 0.555(6)(53)(2) |

**Table 3:** Comparison of the $B's$ at 3 GeV in the SUSY basis in the $\overline{\text{MS}}$ scheme of [66]. *We do not recommend to use the bag as a parametrisation of the BSM four-quark operators, we only quote them in order to compare the results obtained by different collaborations.* In ETM′12 and RBC – UKQCD′12 the renormalisation was performed in the RI – MOM scheme with exceptional kinematics whereas in SWME′14 it is performed perturbatively. The new preliminary RBC-UKQCD results are computed using different intermediate schemes and clearly show that the intermediate scheme difference is much larger than expected (an $\alpha_s^2$ effect). We argue that the renormalisation procedure is the cause of the disagreement observed for $B_4$ and $B_5$ between the different collaborations and that it is due to some underestimated systematic errors present in the $RI-MOM$ scheme. See the text for more detailed explanation. The errors have been already combined, except for SWME'15 where the first errors are statistical and the second systematic. For the RBC-UKQCD'15, the errors are statistics, discretisation and chiral respectively. All the other errors are already combined.

ogy . The $\Delta I = 1/2$ puzzle seems to be explained by the non-perturbative effects [47]. Regarding indirect CP violation, $B_K$ is now known with an impressive precision. The various investigations of the $\Delta S = 2$ BSM operators are converging, the discrepancies observed by several collaborations are likely to be due to systematic errors affecting the non-perturbative renormalisation procedure in RI-MOM schemes. Although a careful study is required, the solution could be provided by the SMOM schemes, which have a much better behaviour. Future improvements will also require to match the lattice computation to phenomenology at a much higher scale in order to decrease the error due to perturbation theory. I have presented here the new determinations of $K \to \pi\pi$ decay amplitudes and neutral kaon mixing matrix elements, but there are other new interesting developments that I have not mentioned here, such as rare kaon decys and the $K_L - K_s$ mass differrence (see [78] and [79] ).

**Acknowledgements-** I would like to thank the organisers for such an enjoyable conference and for promoting my talk. In particular, the conveners of the Goldstone boson working group, Mario Antonelli, Sebastien Descotes-Genon, Andreas Jüttner, Emilie Passemar and Michele Viviani, not only for the organisation, but also the physics discussion we had. I am indebted to Lewis Tunstall for discussing his work on Kaon decay and Chiral perturbation theory. I also would like to thank my colleagues of the RBC-UKQCD collaboration and my lattice colleagues who shared their work with me, sometimes prior to publication.

## References


[1] Z. Bai et al., (2015), 1505.07863.

[2] C. DeTar, (2015), 1511.06884.

[3] M. Della Morte, Proceedings, 50th Recontres de Moriond Electroweak interactions and unified theories, pp. 345–352, 2015, 1507.04051.

[4] L. Lellouch, (2011) 629, 1104.5484.

[5] J.H. Christenson et al., Phys. Rev. Lett. 13 (1964) 138.

[6] NA48, V. Fanti et al., Phys. Lett. B465 (1999) 335, hep-ex/9909022.

[7] KTeV, A. Alavi-Harati et al., Phys. Rev. Lett. 83 (1999) 22, hep-ex/9905060.

[8] G. Buchalla, A.J. Buras and M.E. Lautenbacher, Rev. Mod. Phys. 68 (1996) 1125, hep-ph/9512380.

[9] V. Cirigliano et al., Rev. Mod. Phys. 84 (2012) 399, 1107.6001.

[10] M. Constantinou et al., PoS LATTICE2014 (2014) 390, 1412.1351.

[11] RBC Collaboration, T. Blum et al., Phys.Rev. D68 (2003) 114506, hep-lat/0110075.

[12] A.J. Buras, J.M. Gérard and W.A. Bardeen, Eur. Phys. J. C74 (2014) 2871, 1401.1385.

[13] V. Cirigliano et al., Phys. Rev. Lett. 91 (2003) 162001, hep-ph/0307030.

[14] J. Bijnens and A. Celis, Phys. Lett. B680 (2009) 466, 0906.0302.

[15] J. Bijnens and J. Prades, JHEP 06 (2000) 035, hep-ph/0005189.

[16] T. Hambye, S. Peris and E. de Rafael, JHEP 05 (2003) 027, hep-ph/0305104.

[17] W.A. Bardeen, A.J. Buras and J.M. Gerard, Phys. Lett. B192 (1987) 138.

[18] E. de Rafael, Theoretical Advanced Study Institute in Elementary Particle Physics (TASI 94): CP Violation and the limits of the Standard Model Boulder, Colorado, May 29-June 24, 1994, 1995, hep-ph/9502254.

[19] L. Maiani and M. Testa, Phys. Lett. B245 (1990) 585.

[20] C.W. Bernard et al., Phys. Rev. D32 (1985) 2343.

[21] M.B. Gavela et al., Phys. Lett. B211 (1988) 139.

[22] CP-PACS, J.I. Noaki et al., Phys. Rev. D68 (2003) 014501, hep-lat/0108013.

[23] P. Boucaud et al., Nucl. Phys. B721 (2005) 175, hep-lat/0412029.

[24] R.J. Crewther and L.C. Tunstall, 8th International Workshop on Chiral Dynamics (CD 2015) Pisa, Italy, June 29-July 3, 2015, 2015, 1510.01322.

[25] R.J. Crewther and L.C. Tunstall, Phys. Rev. D91 (2015) 034016, 1312.3319.

[26] L. Lellouch and M. Luscher, Commun.Math.Phys. 219 (2001) 31, hep-lat/0003023.

[27] T. Blum et al., Phys.Rev.Lett. 108 (2012) 141601, 1111.1699.

[28] T. Blum et al., Phys. Rev. D86 (2012) 074513, 1206.5142.

[29] RBC, UKQCD, T. Blum et al., (2014), 1411.7017.

[30] G. Martinelli et al., Nucl. Phys. B445 (1995) 81, hep-lat/9411010.

[31] RBC and UKQCD Collaborations, R. Arthur et al., Phys.Rev. D85 (2012) 014501, 1109.1223.

[32] R.C. Brower, H. Neff and K. Orginos, (2012), 1206.5214.

[33] T. Blum et al., Phys. Rev. D91 (2015) 074502, 1502.00263.

[34] T. Blum et al., (2011), 1106.2714.

[35] C.h. Kim and N.H. Christ, Nucl. Phys. Proc. Suppl. 119 (2003) 365, hep-lat/0210003, [,365(2002)].

[36] U.J. Wiese, Nucl. Phys. B375 (1992) 45.

[37] C. Kelly, LATTICE2015 (2015).

[38] A. Jüttner, LATTICE2015 (2015).

[39] N. Ishizuka et al., Phys. Rev. D92 (2015) 074503, 1505.05289.

[40] M.K. Gaillard and B.W. Lee, Phys. Rev. Lett. 33 (1974) 108.

[41] G. Altarelli and L. Maiani, Phys. Lett. B52 (1974) 351.

[42] E. Endress and C. Pena, Phys. Rev. D90 (2014) 094504, 1402.0827.

[43] P. Hernandez et al., JHEP 05 (2008) 043, 0802.3591.

[44] L. Giusti et al., Phys. Rev. Lett. 98 (2007) 082003, hep-ph/0607220.

[45] L. Giusti et al., JHEP 11 (2004) 016, hep-lat/0407007.

[46] C. Pena, S. Sint and A. Vladikas, JHEP 09 (2004) 069, hep-lat/0405028.

[47] RBC, UKQCD, P.A. Boyle et al., Phys. Rev. Lett. 110 (2013) 152001, 1212.1474.

[48] A. Pich and E. de Rafael, Phys. Lett. B374 (1996) 186, hep-ph/9511465.

[49] Y. Aoki et al., Phys.Rev. D84 (2011) 014503, 1012.4178.

[50] S. Durr et al., Phys.Lett. B705 (2011) 477, 1106.3230.

[51] ETM Collaboration, M. Constantinou et al., Phys.Rev. D83 (2011) 014505, 1009.5606.

[52] T. Bae et al., PoS LATTICE2013 (2014) 476, 1310.7319.

[53] RBC Collaboration, UKQCD Collaboration, R. Arthur et al., Phys.Rev. D87 (2013) 094514, 1208.4412.

[54] J. Laiho and R.S. Van de Water, PoS LATTICE2011 (2011) 293, 1112.4861.

[55] SWME, T. Bae et al., Phys. Rev. Lett. 109 (2012) 041601, 1111.5698.

[56] T. Bae et al., Phys. Rev. D82 (2010) 114509, 1008.5179.

[57] C. Aubin, J. Laiho and R.S. Van de Water, Phys. Rev. D81 (2010) 014507, 0905.3947.

[58] RBC, UKQCD, D.J. Antonio et al., Phys. Rev. Lett. 100 (2008) 032001, hep-ph/0702042.

[59] RBC-UKQCD Collaboration, C. Allton et al., Phys.Rev. D78 (2008) 114509, 0804.0473.

[60] M. Papinutto, C. Pena and D. Preti, PoS LATTICE2014 (2014) 281, 1412.1742.

[61] J. Frison, P. Boyle and N. Garron, PoS LATTICE2014 (2015) 285, 1412.0834.

[62] SWME, Y.C. Jang et al., (2015), 1509.00592.

[63] ETM, N. Carrasco et al., Phys. Rev. D92 (2015) 034516, 1505.06639.

[64] C. Allton et al., Phys.Lett. B453 (1999) 30, hep-lat/9806016.

[65] M. Ciuchini et al., JHEP 9810 (1998) 008, hep-ph/9808328.

[66] A.J. Buras, M. Misiak and J. Urban, Nucl.Phys. B586 (2000) 397, hep-ph/0005183.

[67] R. Babich et al., Phys.Rev. D74 (2006) 073009, hep-lat/0605016.

[68] A. Donini et al., Phys.Lett. B470 (1999) 233, hep-lat/9910017.

[69] RBC Collaboration, UKQCD Collaboration, P. Boyle, N. Garron and R. Hudspith, Phys.Rev. D86 (2012) 054028, 1206.5737.

[70] Collaboration for the ETM, V. Bertone et al., JHEP 1303 (2013) 089, 1207.1287.

[71] SWME Collaboration, T. Bae et al., Phys.Rev. D88 (2013) 071503, 1309.2040.

[72] D. Becirevic and G. Villadoro, Phys.Rev. D70 (2004) 094036, hep-lat/0408029.

[73] J.A. Bailey et al., Phys.Rev. D85 (2012) 074507, 1202.1570.

[74] R. Hudspith, N. Garron and A. Lytle, LATTICE2015 (2015).

[75] C. Sturm et al., Phys.Rev. D80 (2009) 014501, 0901.2599.

[76] RBC-UKQCD, A. Lytle et al., PoS LATTICE2013 (2014) 400.

[77] A.J. Buras, Proceedings, 2015 European Physical Society Conference on High Energy Physics (EPS-HEP 2015), 2015, 1510.00128.

[78] RBC, UKQCD, N.H. Christ et al., Phys. Rev. D92 (2015) 094512, 1507.03094.

[79] N.H. Christ et al., Phys. Rev. D91 (2015) 114510, 1504.01170.